

LYNDON FACTORS AND THE FIBONACCI WORD

KALLE SAARI

ABSTRACT. It is a fundamental property of non-letter Lyndon words that they can be expressed as a concatenation of two shorter Lyndon words. This leads to a naive lower bound $\log_2(n) + 1$ for the number of Lyndon factors that a Lyndon word of length n must have. But this bound is not optimal. In this paper we show that a much more accurate lower bound is $\log_\phi(n)$, where $\phi = (1 + \sqrt{5})/2$, the golden ratio. We show that this bound is optimal in that it is attained by the Fibonacci Lyndon words, and that they are the unique Lyndon words doing so. As an application, we introduce a mapping $\mathcal{L}_\mathbf{x}$ that counts the number of Lyndon factors of length at most n of an infinite word $\mathbf{x}$. We show that a recurrent infinite word $\mathbf{x}$ is aperiodic if and only if $\mathcal{L}_\mathbf{x} \geq \mathcal{L}_\mathbf{f}$, where $\mathbf{f}$ is the Fibonacci infinite word, with equality if and only if $\mathbf{x}$ is in the shift orbit closure of $\mathbf{f}$.

Keywords: Lyndon word, Fibonacci word, Central word, Golden ratio, Sturmian word, Periodicity

1. INTRODUCTION

Lyndon words are primitive words that are the lexicographically smallest words in their conjugacy classes [17]. Originally defined in the context of free Lie algebras [6], Lyndon words have shown to be a useful tool for a variety of problems in combinatorics ranging from the construction of de Bruijn sequences [15] to proving the optimal lower bound for the size of uniform unavoidable sets [5]. One of the fundamental properties of Lyndon words is their recursive nature: if w is a non-letter Lyndon word, then there exist two shorter Lyndon words u and v such that $w = uv$. This implies that the number of different Lyndon factors of w is bounded below by $\log_2|w| + 1$, but a little experimentation quickly shows that this is hardly optimal. One of the results of this paper (Corollary 1) is that a much better lower bound is $\log_\phi|w|$, where $\phi = (1 + \sqrt{5})/2$ is the golden ratio. This bound is optimal in the sense that the Fibonacci Lyndon words attain it. In fact, they are the unique words doing so. More precisely (Theorem 1), we show that if w is a Lyndon word with $|w| \geq F_n$, where F_n is the n^{th} Fibonacci number, then the number of Lyndon factors in w is at least n with equality if and only if w equals one of the two Fibonacci Lyndon words of length F_n , up to renaming the letters.

It also makes sense to count the number of Lyndon factors of infinite words, but there we have to use caution: if an infinite word is aperiodic, it will have infinitely many Lyndon factors (Corollary 2). Thus we define a mapping $\mathcal{L}_\mathbf{x}: \mathbb{N} \rightarrow \mathbb{N}$ for which $\mathcal{L}_\mathbf{x}(n)$ is the number of distinct Lyndon factors of length at most n of a given infinite word $\mathbf{x}$. Of special importance is the Fibonacci infinite word $\mathbf{f}$. Our first main result (Theorem 3) in this setting is that if $\mathbf{x}$ is aperiodic, then $\mathcal{L}_\mathbf{x} \geq \mathcal{L}_\mathbf{f}$. This is a significant improvement of a classic result by Ehrenfeucht and Silberger [12] who showed that an aperiodic infinite word must have arbitrarily long unbordered factors. Indeed, Lyndon words being unbordered, we can now say exactly how many unbordered factors of at most a given length there must

be. If we confine ourselves to recurrent infinite words, which is not a serious restriction, then the above result can be improved as follows (Theorem 4). A recurrent infinite word $\mathbf{x}$ is aperiodic if and only if $\mathcal{L}_{\mathbf{x}} \geq \mathcal{L}_{\mathbf{f}}$ with equality if and only if $\mathbf{x}$ is in the shift orbit closure of the Fibonacci word $\mathbf{f}$, to renaming the letters.

The Fibonacci word is a kind of a universal optimality prover in that it possesses a wide range of extremal properties, see e.g. [3, 7, 10, 19].

2. PRELIMINARIES

In this section we establish the notation of this paper and present some preliminary results. We assume the reader is familiar with the usual terminology of words and languages as given in [1] or [18].

Let $\mathcal{A}$ be a finite, nonsingular alphabet totally ordered by $<$; thus every pair of distinct letters $a, b \in \mathcal{A}$ satisfy either $a < b$ or $b < a$, but not both. We use the same symbol ' $<$ ' to denote the usual order relation among the integers, but this should not cause problems as the context always tells which order is meant. In what follows, we sometimes assume that $0, 1 \in \mathcal{A}$, sometimes $a, b \in \mathcal{A}$, and then their mutual order is implicitly assumed to be their “natural order,” so that $0 < 1$ and $a < b$.

The set of all finite words over $\mathcal{A}$ is denoted by $\mathcal{A}^*$ and the set of finite words excluding the empty word ε is denoted by $\mathcal{A}^+$.

Let $w = a_1 a_2 \cdots a_n$ be a nonempty finite word with $a_i \in \mathcal{A}$ and $n \geq 1$. The *length* of w is $|w| = n$. (We denote the cardinality of a set X by $\#X$.) The *reversal* of w is the word $w^R = a_n a_{n-1} \cdots a_1$. If $w^R = w$, then w is a *palindrome*. The word w has period $p \geq 1$ if $a_{i+p} = a_i$ for all $i = 1, 2, \dots, n - p$. According to this definition, any $p \geq n$ is a period of w . If $p \leq n$, then p is a period of w if and only if there exist words $x, y, z \in \mathcal{A}^*$ such that $w = xy = zx$ and $|y| = |z| = p$. If w has no periods smaller than $|w|$, then it is called *unbordered*, otherwise w is *bordered*. Suppose that $w = uvt$ with $u, v, t \in \mathcal{A}^*$. Then u, v, t are called a *prefix*, *factor*, and *suffix* of w , respectively. In addition, u and t are *proper* if they do not equal w . We say that a word $z \in \mathcal{A}^+$ is a *periodic extension* of w if z is a prefix of a word in w^+ . We abuse the word “extension” here in that we allow an “extension” to be a prefix of w . The word we get from w by deleting its last letter is denoted by w^b ; thus $w^b = a_1 a_2 \cdots a_{n-1}$. Also if $w = xy$ for some words x, y , we denote $x^{-1}w = y$ and $wy^{-1} = x$. The word w is *primitive* if it cannot be written in the form $w = u^k$ for a word $u \in \mathcal{A}^+$ and an integer $k \geq 2$. If $w = uv$, then the word vu is called a *conjugate* of w . The set of all conjugates of w is called the *conjugacy class* of w .

Lemma 1 (Castelli, Mignosi, and Restivo [4]). Let $w \in \mathcal{A}^+$ be a word with periods p, q .

- (i) If $q < p \leq |w|$, then the prefix and suffix of w of length $|w| - q$ have periods q and $p - q$.
- (ii) Let u and v be the prefix and suffix of w of length q , respectively. Then uw and wv have periods q and $p + q$.

In the property (ii) in the previous lemma, the indicated source [4] only mentions and proves the claim for the periods of uw , but the case for the periods of wv can be proved similarly.

Lemma 2 (Fine and Wilf [14]). If a word $w \in \mathcal{A}^+$ has periods p and q , and

$$p + q - \gcd(p, q) \leq |w|,$$

then w has period $\gcd(p, q)$.

It is well-known that the above lemma is optimal. That is to say, if neither of p and q equals $\gcd(p, q)$, then there exists a word $z \in \mathcal{A}^+$ of length $|z| = p + q - \gcd(p, q) - 1$ with periods p and q , but no period $\gcd(p, q)$. Following [18], we call a word $z \in \{a, b\}^*$ a *central word* over $\{a, b\}$ if there exist two coprime integers p, q such that $|z| = p + q - 2$ and both p and q are periods of z . Equivalently, z is a central word over $\{a, b\}$ if either $z \in a^* \cup b^*$ or there exist two coprime integers $p, q \geq 2$ such that $|z| = p + q - 2$, both p and q are periods of z , but z does not have period $\gcd(p, q) = 1$, that is, both letters a and b occur in z . These words are also known as extremal Fine and Wilf words [21]. They are palindromes and unique up to letter-to-letter isomorphism [18, 21]. The latter fact implies that there are exactly two central words over $\{a, b\}$ with given periods p and q ; if one is z , then the other one is $c(z)$, where c is the morphism $a \mapsto b, b \mapsto a$.

Recall that the Fibonacci numbers are defined recursively as $F_1 = 1, F_2 = 1$, and $F_n = F_{n-1} + F_{n-2}$ for $n \geq 3$, and that every two consecutive Fibonacci numbers are coprime. The two central words over $\{a, b\}$ for periods F_{n-2} and F_{n-1} can be obtained as follows. Let $a, b \in \mathcal{A}$ be distinct letters. Then define $f_1 = b, f_2 = a$, and

$$f_n = f_{n-1}f_{n-2} \quad (n \geq 3).$$

We call the words f_n *finite Fibonacci words* over $\{a, b\}$. Note that we could also have defined $f_1 = a$ and $f_2 = b$. We will make this distinction when it matters. Let p_n denote the word for which $f_n = p_n xy$ with $xy \in \{ab, ba\}$ and $n \geq 3$. Thus

$$\begin{array}{lllll} f_3 = ab & f_4 = aba & f_5 = abaab & f_6 = abaabaab & f_7 = abaababaabaab \\ p_3 = \varepsilon & p_4 = a & p_5 = aba & p_6 = abaaba & p_7 = abaababaaba \end{array}$$

Note that $|f_n| = F_n$ and thus $|p_n| = F_n - 2$. It can be shown that if $n \geq 5$, then p_n has periods F_{n-2} and F_{n-1} , but it does not have period $\gcd(F_{n-2}, F_{n-1}) = 1$. Thus each p_n is a central word over $\{a, b\}$. The other central word with the same periods and is $c(p_n)$, where c is the morphism $a \mapsto b$ and $b \mapsto a$.

The order $<$ of $\mathcal{A}$ is extended to $\mathcal{A}^*$ as follows: For $u, v \in \mathcal{A}^*$ we have

$$u < v \iff \begin{cases} u \text{ is a prefix of } v, \text{ or} \\ u = xau' \text{ and } v = xbv' \text{ with } x, u', v' \in \mathcal{A}^*, a, b \in \mathcal{A} \text{ and } a < b. \end{cases}$$

This is called the *lexicographic ordering* of $\mathcal{A}^*$ with respect to $<$.

A nonempty, primitive word $w \in \mathcal{A}^+$ is called a *Lyndon word* if it is the smallest word in its conjugacy class. In particular, letters are Lyndon words, but the empty word is not. For example, the Lyndon words $w \in \{0, 1\}^+$ with $|w| \leq 4$ are

$$0, \quad 1, \quad 01, \quad 001, \quad 011, \quad 0001, \quad 0011, \quad 0111.$$

Lemma 3 (Berstel and de Luca [2]). If $z \in \mathcal{A}^*$ is a central word over $\{a, b\}$ and $a < b$, then azb is a Lyndon word.

According to Lemma 3, the words ap_nb and $ac(p_n)b$ are Lyndon words; we call them *Fibonacci Lyndon words of length F_n over $\{a, b\}$* . The first few ones are

$$\begin{array}{lllll} ab & aab & aabab & aabaabab & aabaababaabab \\ ab & abb & ababb & ababbabb & ababbababbabb \end{array}$$

The properties of Lyndon words given in the next lemma are well-known [17, 11].

Lemma 4. Lyndon words have following properties.

- (i) Lyndon words are unbordered.
- (ii) A word w is Lyndon if and only if $w < y$ for all nonempty suffixes y of w .

(iii) If u and v are Lyndon words and $u < v$, then uv is a Lyndon word.

Lemma 5. Let $w \in \mathcal{A}^+$ be a Lyndon word. Suppose that za is a periodic extension of w for some $z \in \mathcal{A}^+$ and $a, b \in \mathcal{A}$ with $a < b$. Then zb is a Lyndon word.

Proof. There exist an integer $n \geq 0$ and words $x, y \in \mathcal{A}^*$ such that $w = xay$ and $z = w^n x$. We show that xb is a Lyndon word; this suffices because then Lemma 4 implies that $zb = w^n xb$ is a Lyndon word since $w < xb$.

Contrary to what we want to show, suppose that xb is not a Lyndon word. Then Lemma 4 implies that xb has a nonempty suffix v with $v < xb$. Write $v = v'b$. Then $v'ay$ is a suffix of w , so that $v'ay > w$ because w is a Lyndon word. Therefore $b > a$ implies that $v = v'b$ is not a prefix of w and thus not a prefix of x . Consequently $v < xb$ implies that we can write $v = tct'$ and $xb = tdt''$ for some words $t, t', t'' \in \mathcal{A}^*$ and letters $c, d \in \mathcal{A}$ with $c < d$. Since $|td| = |tc| \leq |v| < |xb|$, the word td is a prefix of x and thus a prefix of w . But then

$$v'ay < v'by = vy = tct'y < td < w,$$

a contradiction. Thus xb is a Lyndon word, and the proof is complete. $\square$

Lemma 6. Let $w \in \mathcal{A}^+$ be a Lyndon word with $|w| \geq 2$. Let λ_w be the longest proper prefix of w that is also a Lyndon word. Then w^b is a periodic extension of λ_w . Furthermore, the word $\mu_w := \lambda_w^{-1}w$ is a Lyndon word.

Proof. Suppose that w^b is not a periodic extension of λ_w . Then there exist a word u and different letters a, b such that ua is a prefix of λ_w and $\lambda_w^k ub$ is a prefix of w^b for some integer $k \geq 1$. If $a < b$, then $\lambda_w^k ub$ is a Lyndon word by Lemma 5, contradicting the maximality of $|\lambda_w|$. If $a > b$, then $(\lambda_w^k)^{-1}w < w$ because $(\lambda_w^k)^{-1}w$ begins with ub while w begins with ua . Thus w has a nonempty suffix that is smaller than w , contradicting Lemma 4 because w is Lyndon. Therefore w^b is a periodic extension of λ_w , and consequently there exist $u \in \mathcal{A}^+$ and different letters $a, b \in \mathcal{A}$ such that ua is a prefix of λ_w and $w = \lambda_w^k ub$ for some integer $k \geq 1$. Furthermore, $a < b$ because otherwise w would not be a Lyndon word. Now $\mu_w = \lambda_w^{-1}w = \lambda_w^{k-1}ub$ is a Lyndon word by Lemma 5. $\square$

Due to its importance in upcoming considerations, let us restate Lemma 6: every non-letter Lyndon word $w \in \mathcal{A}^+$ can be written as $w = \lambda_w \mu_w$, where λ_w and μ_w are Lyndon words and w^b is a periodic extension of λ_w .

An *infinite word* is a sequence $\mathbf{x} = a_1 a_2 a_3 \dots a_n \dots$ where $a_n \in \mathcal{A}$. The set of infinite words over $\mathcal{A}$ is denoted by $\mathcal{A}^{\mathbb{N}}$. A *tail* of the infinite word $\mathbf{x}$ is another infinite word $\mathbf{y} \in \mathcal{A}^{\mathbb{N}}$ such that $\mathbf{x} = x\mathbf{y}$ for some $x \in \mathcal{A}^*$. An infinite word $\mathbf{x}$ is *purely periodic* if $\mathbf{x} = uuu \dots u \dots$ for some finite word $u \in \mathcal{A}^+$; we also denote this by $\mathbf{x} = u^\omega$. The word $\mathbf{x}$ is *ultimately periodic* if it has a purely periodic tail. Finally, $\mathbf{x}$ is *aperiodic* if it is not ultimately periodic. A *factor* of $\mathbf{x}$ is a finite word that occurs somewhere in $\mathbf{x}$. The set of all factors of $\mathbf{x}$ is denoted by $F(\mathbf{x})$. The word $\mathbf{x}$ is called *recurrent* if each of its factors occurs at least twice (and thus infinitely many times) in $\mathbf{x}$. If an infinite word is ultimately periodic and recurrent, then it is actually purely periodic. The *shift orbit closure* of $\mathbf{x}$ is the set of all infinite words $\mathbf{y} \in \mathcal{A}^{\mathbb{N}}$ such that $F(\mathbf{y}) \subseteq F(\mathbf{x})$. We denote the Lyndon factors of $\mathbf{x}$ by $L(\mathbf{x})$. (Do not confuse this with the function $\mathcal{L}_x$ defined in Section 4.)

Let f_n be the finite Fibonacci words over $\{a, b\}$ defined above with $f_1 = b$ and $f_2 = a$. If $n \geq 2$, then f_n is a prefix of f_{n+1} . Thus there exists a unique infinite word $\mathbf{f}$ such that f_n is a prefix of $\mathbf{f}$ for every $n \geq 2$. The word $\mathbf{f}$ is called a *Fibonacci infinite word* over $\{a, b\}$. Note that there is another Fibonacci infinite word over $\{a, b\}$, which results from

defining $f_1 = a$ and $f_2 = b$. When we want to stress the definition of f_1 and f_2 when constructing $\mathbf{f}$, we denote $\mathbf{f} = \lim_{n \rightarrow \infty} f_n$. The infinite Fibonacci words are an archetype of Sturmian words [18]. In particular, they are recurrent.

Lemma 7. The Fibonacci word $\mathbf{f} = \lim_{n \rightarrow \infty} f_n$ has the following properties.

- (1) The words p_n are palindromes [9].
- (2) The Lyndon factors of $\mathbf{f}$ are precisely the Lyndon conjugates of f_n [8, Lemma 7].
- (3) If w is a conjugate of f_n , then the reversal w^R is a conjugate of f_n [22].

Lemma 8. Let $\mathbf{f} = \lim_{n \rightarrow \infty} f_n$ be a Fibonacci word for which $\{f_1, f_2\} = \{a, b\}$. Write $f_n = p_n xy$ with $xy \in \{ab, ba\}$ and suppose that $a < b$. Then the Lyndon conjugate of f_n is the word ap_nb for all $n \geq 3$. Furthermore, every Lyndon factor of $\mathbf{f}$ that is shorter than ap_nb is either a prefix or a suffix of ap_nb .

Proof. First off, the word ap_nb is a Lyndon word by Lemma 3. Thus the first claim is proved by showing that ap_nb is a conjugate of f_n . This is clear if $f_n = p_n ba$, so assume that $f_n = p_n ab$ instead. Then bp_na is a conjugate of f_n . Since the reversal of a conjugate of f_n is a conjugate of f_n and since p_n is a palindrome by Lemma 7, it follows that $ap_n^R b = ap_nb$ is a conjugate of f_n .

Next we show that if $k < n$, then the Lyndon conjugate of f_k is a prefix or a suffix of ap_nb . Since $\{f_1, f_2\} = \{a, b\}$, the claim is plainly true for $k = 1, 2$. Also, a quick case analysis shows that the Lyndon conjugate of f_3 , which is ab , is either a prefix or suffix of ap_nb . Thus we may suppose that $k \geq 4$. Then $k < n$ implies that f_k is a prefix of p_n . Furthermore, since p_n is a palindrome, the reversal f_k^R is a suffix of p_n . Therefore, if $f_k = p_k ba$, then $ap_k b$ is a prefix of ap_nb ; if $f_k = p_k ab$, then $ap_k b$ is a suffix of ap_nb . $\square$

3. LYNDON FACTORS OF LYNDON WORDS

Let $w \in \mathcal{A}^+$ be a Lyndon word. We denote the number of distinct Lyndon factors of w by $\mathcal{L}(w)$. A trivial but useful observation is that if $|w| \geq 2$, then

$$\mathcal{L}(w) \geq \mathcal{L}(\lambda_w) + 1 \quad \text{and} \quad \mathcal{L}(w) \geq \mathcal{L}(\mu_w) + 1,$$

where λ_w and μ_w are the Lyndon words provided by Lemma 6. If $|w| \geq 2$, let p_w denote the word such that $w = ap_w b$ for some letters $a, b \in \mathcal{A}$.

Lemma 9. If $w \in \mathcal{A}^+$ is a Fibonacci Lyndon word of length F_n with $n \geq 3$, then $\mathcal{L}(w) = n$.

Proof. This is an immediate corollary to Lemma 8: the word w is the Lyndon conjugate of f_n (for some choice of $f_1, f_2 \in \mathcal{A}$). Each of the Lyndon conjugates of f_k with $1 \leq k \leq n$ are either prefixes or suffixes of w , and these are the only Lyndon factors of w . Thus $\mathcal{L}(w) = n$. $\square$

Lemma 10. Let $w \in \mathcal{A}^+$ be a Lyndon word with $|w| = F_n$ for $n \geq 3$ and let $a, b \in \mathcal{A}$ be the letters such that $w = ap_w b$. Then w is a Fibonacci Lyndon word over $\{a, b\}$ if and only if ap_w has period F_{n-1} and $p_w b$ has period F_{n-2} , or vice versa.

Proof. Suppose that w is a Fibonacci Lyndon word over $\{a, b\}$. The claim is readily verified for $n \leq 4$, so assume that $n \geq 5$. Since p_w has periods F_{n-2} and F_{n-1} but does not have period $\gcd(F_{n-2}, F_{n-1}) = 1$, it follows that the word ap_w has period either F_{n-2} or F_{n-1} . Indeed, if p_w did not have either period, then bp_w would have both periods, contradicting Lemma 2 because $|p_w| = F_n - 2$. An analogous argument shows that $p_w b$

must have period either F_{n-2} or F_{n-1} . Finally, ap_w and $p_w b$ cannot have the same period F_{n-2} or F_{n-1} because otherwise w would have the same period, which it does not since it is unbordered by Lemma 4.

Conversely, suppose that ap_w has period F_{n-1} and $p_w b$ has period F_{n-2} , or vice versa. Again if $n \leq 4$, the claim is trivial, so assume $n \geq 5$. Then since $F_{n-1} < |ap_w| = |p_w b|$, it follows that both a and b occur in p_w , and thus $\gcd(F_{n-1}, F_{n-2}) = 1$ is not a period of p_w . Since p_w also satisfies $|p_w| = F_n - 2$ and it has periods F_{n-2} and F_{n-1} , we have shown that p_w is a central word and thus w is a Fibonacci Lyndon word over $\{a, b\}$. $\square$

Lemma 11. Let $w \in \mathcal{A}^+$ be a Lyndon word with $|w| \geq F_n$ for some $n \geq 3$. We have $\mathcal{L}(w) \geq n$. Furthermore, the identity $\mathcal{L}(w) = n$ implies that w is a Fibonacci Lyndon word of length F_n .

Proof. Using the fact that letters are Lyndon words and that w is a product of two shorter Lyndon words, the reader readily verifies the claim for $n \leq 4$. Hence we assume that $n \geq 5$. Since w is a Lyndon word, we have $w = \lambda_w \mu_w = ap_w b$ for some letters $a, b \in \mathcal{A}$ with $a < b$. We split the proof in several cases depending on the length of λ_w .

Case (i). Suppose that $|\lambda_w| > F_{n-1}$. Since $n - 1 \geq 4$, we may apply the induction assumption to λ_w and obtain

$$\mathcal{L}(w) \geq \mathcal{L}(\lambda_w) + 1 \geq n + 1 > n.$$

Case (ii). Suppose that $|\lambda_w| = F_{n-1}$. Then $|\mu_w| \geq F_{n-2}$, and we have three subcases:

Case (ii-a). If μ_w is not a factor of λ_w , then the induction assumption implies

$$\mathcal{L}(w) \geq \mathcal{L}(\lambda_w) + \#\{w, \mu_w\} \geq (n - 1) + 2 > n.$$

Case (ii-b). Suppose that μ_w is a factor but not a suffix of λ_w . Then by denoting the longest Lyndon prefix of λ_w by λ_{λ_w} , we have $|\lambda_{\lambda_w}| \geq |\mu_w|$ because λ_w^b is a periodic extension of λ_{λ_w} by Lemma 6 and μ_w is unbordered by Lemma 4. Therefore $|\lambda_{\lambda_w}| \geq F_{n-2}$. Since $n - 2 \geq 3$, we may apply the induction assumption to λ_{λ_w} , obtaining

$$(1) \quad \mathcal{L}(w) \geq \mathcal{L}(\lambda_w) + 1 \geq \mathcal{L}(\lambda_{\lambda_w}) + 2 \geq (n - 2) + 2 = n,$$

where the last inequality is equality only if $|\mu_w| = |\lambda_{\lambda_w}| = F_{n-2}$. This would imply that λ_{λ_w} and μ_w are conjugates, which would further imply that $\lambda_{\lambda_w} = \mu_w$ because both are Lyndon words. But then μ_w is both a prefix and a suffix of w , contradicting the fact that w is unbordered. Hence the third inequality in (1) is strict.

Case (ii-c). Suppose that μ_w is a suffix of λ_w . Then it is actually a suffix of $p_{\lambda_w} b$ because μ_w cannot equal λ_w . Since $|\lambda_w| = F_{n-1}$, the induction assumption gives

$$\mathcal{L}(w) \geq \mathcal{L}(\lambda_w) + 1 \geq (n - 1) + 1 = n.$$

If $\mathcal{L}(w) > n$, we are done, so assume $\mathcal{L}(w) = n$. Then $\mathcal{L}(\lambda_w) = n - 1$. We will show that ap_w has period F_{n-1} and $p_w b$ period F_{n-2} , which means that w is a Fibonacci Lyndon word by Lemma 10.

First, the word ap_w has period $|\lambda_w| = F_{n-1}$ because $ap_w = w^b$ is a periodic extension of λ_w by Lemma 6. Second, since $|\lambda_w| = F_{n-1}$ and $\mathcal{L}(\lambda_w) = n - 1$, the induction assumption implies that λ_w is a Fibonacci Lyndon word and therefore Lemma 6 implies that $p_{\lambda_w} b$ has period either F_{n-3} or F_{n-2} . The period cannot be F_{n-3} , however, because the unbordered word μ_w is a suffix of $p_{\lambda_w} b$ and $|\mu_w| \geq F_{n-2} > F_{n-3}$. Thus $p_{\lambda_w} b$ has period F_{n-2} , and furthermore $|\mu_w| = F_{n-2}$. Now the fact that μ_w is a suffix of $p_{\lambda_w} b$ with $|\mu_w| = F_{n-2}$ and that $p_{\lambda_w} b$ has period F_{n-2} imply that $p_w b = p_{\lambda_w} b \mu_w$ has period F_{n-2} by Lemma 1.

Case (iii). Suppose that $|w|/2 < |\lambda_w| < F_{n-1}$. Then $|\mu_w| = |w| - |\lambda_w|$ gives $F_{n-2} < |\mu_w| < |\lambda_w|$, so λ_w is not a factor of μ_w . Furthermore, since $n - 2 \geq 3$, the induction assumption gives

$$\mathcal{L}(w) \geq \mathcal{L}(\mu_w) + \#\{w, \lambda_w\} > (n - 2) + 2 = n.$$

Case (iv). Suppose that $|\lambda_w| = |w|/2$. Then $|\mu_w| = |\lambda_w|$ and thus μ_w is not a factor of λ_w because otherwise $\mu_w = \lambda_w$ and w would be bordered. Noting that $|\lambda_w| = |w|/2 > F_{n-2}$ because $n \geq 4$, we therefore have by induction

$$\mathcal{L}(w) \geq \mathcal{L}(\lambda_w) + \#\{w, \mu_w\} > (n - 2) + 2 = n.$$

Case (v). Suppose that $F_{n-2} < |\lambda_w| < |w|/2$. Then $|\lambda_w| < |\mu_w|$, and so μ_w is not a factor of λ_w . Since we also have $n - 2 \geq 3$, the induction assumption gives

$$\mathcal{L}(w) \geq \mathcal{L}(\lambda_w) + \#\{w, \mu_w\} > (n - 2) + 2 = n.$$

Case (vi). Suppose that $|\lambda_w| = F_{n-2}$. Then $|\mu_w| \geq F_{n-1}$, so the induction assumption gives

$$(2) \quad \mathcal{L}(w) \geq \mathcal{L}(\mu_w) + 1 \geq (n - 1) + 1 = n.$$

If $\mathcal{L}(w) > n$, we are done, so assume $\mathcal{L}(w) = n$. Our goal is to show that ap_w has period F_{n-2} and $p_w b$ has period F_{n-1} , which means that w is a Fibonacci Lyndon word by Lemma 10. The first objective is easy because $w^b = ap_w$ is a periodic extension of λ_w by Lemma 6, and thus ap_w has period $|\lambda_w| = F_{n-2}$.

For the second objective, we take care of a special case first. If $n = 5$, then $|\lambda_w| = 2$, so that $\lambda_w = ab$. Then $w = ababb$ because w^b is a periodic extension of λ_w , and consequently $p_w b = 1011$ has period F_4 , as claimed. We may thus assume that $n \geq 6$.

Now, note that since $\mathcal{L}(w) = n$ in Eq. (2), we have $\mathcal{L}(\mu_w) = n - 1$. Since also $|\mu_w| \geq F_{n-1}$, the induction assumption says that actually $|\mu_w| = F_{n-1}$ and that μ_w is a Fibonacci Lyndon word. Thus Lemma 6 implies that one of ap_{μ_w} and $p_{\mu_w} b$ has period F_{n-3} and the other one period F_{n-2} . Since $w^b = \lambda_w ap_{\mu_w}$ is a periodic extension of λ_w and $|\lambda_w| < |\mu_w|$, we see that λ_w is a prefix of ap_{μ_w} . Therefore ap_{μ_w} cannot have period F_{n-3} because $F_{n-3} < |\lambda_w|$ and λ_w is unbordered. Hence ap_{μ_w} has period F_{n-2} and $p_{\mu_w} b$ has period F_{n-3} . Since $n \geq 6$, we have $|p_{\lambda_w} ba| = F_{n-2} \leq F_{n-1} - 2 = |p_{\mu_w}|$, and consequently since $p_w = p_{\lambda_w} ba p_{\mu_w}$ and p_w has period F_{n-2} , it follows that $p_{\lambda_w} ba$ is a prefix of p_{μ_w} . Since p_{μ_w} has periods F_{n-3} and F_{n-2} , Lemma 1 implies that p_w has periods F_{n-2} and $F_{n-2} + F_{n-3} = F_{n-1}$. Now, as we have reasoned before, $p_w b$ must have period either F_{n-2} or F_{n-1} for otherwise $p_w a$ would have both periods contradicting Lemma 2. Since ap_w has period F_{n-2} and w is unbordered, we conclude that $p_w b$ must have period F_{n-1} .

Case (vii). Suppose that $|\lambda_w| < F_{n-2}$. Then $|\mu_w| > F_{n-1}$, so that the induction assumption implies

$$\mathcal{L}(w) \geq \mathcal{L}(\mu_w) + 1 > (n - 1) + 1 = n.$$

□

Theorem 1. Let $w \in \mathcal{A}^+$ be a Lyndon word with $|w| \geq F_n$ for some $n \geq 3$. Then $\mathcal{L}(w) \geq n$ with equality if and only if w is a Fibonacci Lyndon word of length F_n .

Proof. The claim is obtained by combining Lemmas 9 and 11. □

Corollary 1. If $w \in \mathcal{A}^+$ is a Lyndon word, then $\mathcal{L}(w) > \log_\phi |w|$, where $\phi = (1 + \sqrt{5})/2$ is the golden ratio.

Proof. The claim is trivial if $|w| = 1$, so suppose $|w| \geq 2$, and let $n \geq 3$ be the unique integer for which $F_n \leq |w| < F_{n+1}$. It is well-known [16] that $F_{n+1} < \phi^n$, and therefore $\log_\phi |w| < n$. On the other hand, Theorem 1 says that $\mathcal{L}(w) \geq n$. The claim follows from the last two inequalities. $\square$

Remark 1. A noteworthy feature of Theorem 1 is that the optimal words, the Fibonacci Lyndon words, are made of just two different letters. A priori it may seem “obvious” that this should always be the case for a Lyndon word of a given length having the smallest possible number of Lyndon factors, but this, in fact, is not always true. For example, each Lyndon word of length 6 has at least 7 Lyndon factors. In this case the optimal words are, up to renaming the letters,

000001 000101 001101 010111 010102 010202 021022 011111,

three of which are made of three different letters. However, see Conjecture 1.

Conjecture 1. For $n \geq 1$, denote

$$\ell(n) = \min \{ \mathcal{L}(w) \mid w \text{ is a Lyndon word with } |w| = n \}.$$

If w is a Lyndon word with $|w| \neq 6$ and $\mathcal{L}(w) = \ell(|w|)$, then w is Sturmian Lyndon word, i.e., we have $w \in \{a, b\}^+$, $w = ap_w b$, and p_w is a central word. Furthermore, we have

$$\ell_n = \min \left\{ \sum_{i=0}^k a_i \mid \text{there exists an integer } q \text{ such that } n/q = [a_0; a_1, \dots, a_k] \right\}.$$

The expression $[a_0; a_1, \dots, a_k]$ above denotes the continued fraction expansion of n/q .

4. LYNDON FACTORS OF RECURRENT WORDS

Our main goal in this section is to prove Theorem 4 stating that the number of Lyndon factors in recurrent infinite words is governed by the Fibonacci word. We begin with results that are interesting in their own right.

Lemma 12 (Siromoney, Mathew, Dare, and Subramanian [20]). Each infinite word $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$ admits a unique factorization of the form either

$$\mathbf{x} = \prod_{i \geq 1} w_i \quad \text{or} \quad \mathbf{x} = w_1 w_2 \cdots w_n \mathbf{x}',$$

where each $w_i \in \mathcal{A}^+$ is a finite Lyndon word with $w_i \geq w_{i+1}$ and $\mathbf{x}' \in \mathcal{A}^{\mathbb{N}}$ begins with arbitrarily long Lyndon words.

Lemma 13. Let $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$ be an infinite word. Then at least one of the following holds:

- (i) $\mathbf{x}$ is ultimately periodic;
- (ii) $\mathbf{x}$ has a tail that begins with arbitrarily long Lyndon words;
- (iii) $\mathbf{x} = \prod_{i \geq 1} w_i$, where each $w_i \in \mathcal{A}^+$ is a Lyndon word and, for every $k \geq 1$, there exists an index i_k such that $|w_i| > k$ for all $i > i_k$.

Proof. Let us suppose that $\mathbf{x}$ is aperiodic and that it does not have a tail beginning with arbitrarily long Lyndon words; we will show that then $\mathbf{x}$ satisfies property (iii). Let $k \geq 1$. Lemma 12 says that $\mathbf{x}$ admits a factorization $\mathbf{x} = \prod_{i \geq 1} w_i$ in which the $w_i \in \mathcal{A}^+$ are Lyndon words and $w_i \geq w_{i+1}$. Since $\mathbf{x}$ is aperiodic, the sequence of words w_i is not ultimately constant. Therefore we can define

$$i_k = \max \{ j : w_j = \min \{ w_i : |w_i| \leq k \} \}.$$

In other words, i_k is the largest index such that w_{i_k} is the least word among the words w_i of length $|w_i| \leq k$ (there are only finitely many of them because $\mathcal{A}$ is finite). Then $|w_i| > k$ whenever $i > i_k$. Indeed, if $i > i_k$, then the inequality $w_{i_k} \geq w_i$ and the maximality of i_k imply $w_{i_k} > w_i$. Because of the minimality of w_{i_k} , we thus have $|w_i| > k$. $\square$

Corollary 2. If an infinite word $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$ has only finitely many distinct Lyndon factors, then it is ultimately periodic.

Proof. If $\mathbf{x}$ satisfies property (ii) or (iii) in Lemma 13, then it clearly has infinitely many Lyndon factors. Thus $\mathbf{x}$ satisfies property (i) and is ultimately periodic. $\square$

Remark 2. The sleek proof of Corollary 2 was suggested by Tero Harju. The author's original proof was more intricate.

Recall that the set of factors of an infinite word $\mathbf{x}$ is denoted by $F(\mathbf{x})$ and the set of its Lyndon factors is denoted by $L(\mathbf{x})$.

Theorem 2. If $\mathbf{x}$ and $\mathbf{y}$ are recurrent infinite words and $L(\mathbf{x}) = L(\mathbf{y})$, then $F(\mathbf{x}) = F(\mathbf{y})$.

Proof. Suppose first that $\mathbf{x}$ is ultimately periodic. Then, in fact, it is purely periodic because it is recurrent. Writing $\mathbf{x} = u^\omega$, where u is a primitive word, we see that there is only one Lyndon factor of length $|u|$ or more, and that is of course the Lyndon conjugate of u . Thus $\mathbf{y}$ has only finitely many Lyndon factors, so it is ultimately periodic by Corollary 2, and hence purely periodic because it is recurrent. Write $\mathbf{y} = v^\omega$ with v primitive. Since the Lyndon conjugate of u is a factor of $\mathbf{y}$ and the Lyndon conjugate of v is a factor of $\mathbf{x}$, it follows that v and u are conjugates, and thus $F(\mathbf{x}) = F(\mathbf{y})$.

Suppose then that $\mathbf{x}$ is aperiodic. We show that every $u \in F(\mathbf{x})$ is a factor of a Lyndon factor of $\mathbf{x}$. Since $\mathbf{y}$ must be aperiodic as well, the analogous property clearly holds for $\mathbf{y}$, implying that $F(\mathbf{x}) = F(\mathbf{y})$. If $\mathbf{x}$ satisfies property (ii) of Lemma 13, then some tail $\mathbf{x}'$ of $\mathbf{x}$ begins with arbitrarily long Lyndon words. Since $\mathbf{x}$ is recurrent, it follows that u is a factor of a Lyndon prefix of $\mathbf{x}'$. Thus suppose that $\mathbf{x}$ satisfies property (iii) of Lemma 13. Since $\mathbf{x}$ is recurrent, there exists a word v such that $uvu \in F(\mathbf{x})$. Let $k = |uvu|$ and let i_k be the index provided by Lemma 13. Since $\mathbf{x}$ is recurrent, the word uvu occurs in $w_{i_k} w_{i_k+1} w_{i_k+2} \dots$. Since $|w_{i_k+j}| > k$ for each $j \geq 1$, it follows that u necessarily occurs in some w_{i_k+j} . $\square$

For an infinite word $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$, we define a mapping $\mathcal{L}_{\mathbf{x}}: \mathbb{N} \rightarrow \mathbb{N}$ such that $\mathcal{L}_{\mathbf{x}}(n)$ is the number of Lyndon factors of $\mathbf{x}$ of length at most n . Notice that the mapping $\mathcal{L}_{\mathbf{x}}$ is increasing, but not necessarily strictly increasing, as can be seen from Lemma 14. Notice also that $\mathcal{L}_{\mathbf{x}}$ determines the number of Lyndon factors of any length $k \geq 1$. Indeed, it is given by the expression $\mathcal{L}_{\mathbf{x}}(k) - \mathcal{L}_{\mathbf{x}}(k-1)$ for $k \geq 2$.

Lemma 14. Let $\mathbf{f} \in \mathcal{A}^{\mathbb{N}}$ be a Fibonacci infinite word. Then for all $n \geq 1$, we have $\mathcal{L}_{\mathbf{f}}(n) = k$, where $k \geq 2$ is the unique integer such that $F_k \leq n < F_{k+1}$.

Proof. Lemma 8 says that the Lyndon factors in $\mathbf{f}$ are precisely the Lyndon conjugates of the finite Fibonacci words f_k . Therefore if $F_k \leq n < F_{k+1}$, the Lyndon factors of length at most n are the Lyndon conjugates of $f_1, f_2, \dots, f_k$, so that $\mathcal{L}_{\mathbf{f}}(n) = k$. $\square$

Theorem 3. If $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$ is aperiodic, then $\mathcal{L}_{\mathbf{x}} \geq \mathcal{L}_{\mathbf{f}}$.

Proof. Since $\mathcal{L}_{\mathbf{x}}$ is increasing, Lemma 14 implies that it suffices to show that $\mathcal{L}_{\mathbf{x}}(F_k) \geq k$ for all $k \geq 2$. This is clear for $k = 2, 3$ because $\mathbf{x}$ is aperiodic. Thus assume $k \geq 4$, and let w be a shortest Lyndon factor of $\mathbf{x}$ of length $> F_k$; Corollary 2 ensures that such

a word w exists because $\mathbf{x}$ is aperiodic. Theorem 1 implies that $\mathcal{L}(w) > k$. Since w is as short as possible, all of its proper Lyndon factors are of length at most F_k . Therefore

$$\mathcal{L}_{\mathbf{x}}(F_k) \geq \mathcal{L}(w) - 1 \geq k.$$

□

Remark 3. A classic result by Ehrenfeucht and Silberger [12] states that if an infinite word has only finitely many unbordered factors, then it is ultimately periodic. Since Lyndon words are unbordered, Theorem 3 is a quantitative formulation of this with an exact lower bound for the necessary number of unbordered factors.

Remark 4. Looking at Theorem 3, one might be tempted to postulate that if $\mathbf{x}$ is aperiodic, then for all $n \geq 1$, the number of length- n Lyndon factors of $\mathbf{x}$ must be at least as big as the number of length- n Lyndon factors of a Fibonacci infinite word, but this is not true. For example, let $\mathbf{f}$ be the Fibonacci infinite word over $\{a, b\}$ with $f_1 = b$ and $f_2 = a$, and let $\mathbf{x} = g(\mathbf{f})$, where g is the morphism $a \mapsto aab$, $b \mapsto aaab$. Then $\mathbf{x}$ does not have any Lyndon factors of length 5, while $\mathbf{f}$ has the factor $aabab$.

Theorem 4. Let $\mathbf{x} \in \mathcal{A}^{\mathbb{N}}$ be recurrent. Then $\mathbf{x}$ is aperiodic if and only if $\mathcal{L}_{\mathbf{x}} \geq \mathcal{L}_{\mathbf{f}}$ with equality if and only if $\mathbf{x}$ is in the shift orbit closure of a Fibonacci infinite word $\mathbf{f}$.

Proof. Let us first prove the first equivalence. Suppose $\mathbf{x}$ is ultimately periodic. Then it is purely periodic because it is recurrent. Thus $\mathcal{L}_{\mathbf{x}}$ is ultimately constant, so $\mathcal{L}_{\mathbf{x}}(n) < \mathcal{L}_{\mathbf{f}}(n)$ for all sufficiently large n . Conversely, if $\mathcal{L}_{\mathbf{x}}(n) < \mathcal{L}_{\mathbf{f}}(n)$ for some $n \geq 1$, then Theorem 3 implies that $\mathbf{x}$ is ultimately periodic.

Let us next prove the second equivalence. Obviously the condition that $\mathbf{x}$ be in the shift orbit closure of a Fibonacci word $\mathbf{f}$ is sufficient for the identity $\mathcal{L}_{\mathbf{x}} = \mathcal{L}_{\mathbf{f}}$ to hold. Let us prove the converse, and suppose that $\mathcal{L}_{\mathbf{x}} = \mathcal{L}_{\mathbf{f}}$. Then in particular, $\mathcal{L}_{\mathbf{x}}(1) = \mathcal{L}_{\mathbf{f}}(1) = 2$, so that $\mathbf{x}$ consist of two distinct letters, say a and b . Since $\mathcal{L}_{\mathbf{x}}(3) = 1$, exactly one of aab and abb occurs in $\mathbf{x}$. If aab is in $L(\mathbf{x})$, let $\mathbf{f} = \lim_{n \rightarrow \infty} f_n$ be the Fibonacci word with $f_1 = b$ and $f_2 = a$, so that $aab \in L(\mathbf{f})$. If abb is in $L(\mathbf{x})$, let $\mathbf{f} = \lim_{n \rightarrow \infty} f_n$ be the Fibonacci word with $f_1 = a$ and $f_2 = b$, so that $abb \in L(\mathbf{f})$. Then a Lyndon word of length at most 3 is a factor of $\mathbf{x}$ if and only if it is a factor of $\mathbf{f}$. We will show next that $L(\mathbf{x}) = L(\mathbf{f})$; then Theorem 2 implies that $F(\mathbf{x}) = F(\mathbf{f})$ because both $\mathbf{x}$ and $\mathbf{f}$ are recurrent.

If $L(\mathbf{x}) \neq L(\mathbf{f})$, then the identity $\mathcal{L}_{\mathbf{x}} = \mathcal{L}_{\mathbf{f}}$ implies that there exist an integer and Lyndon words w, z with $|w| = |z| = F_k$ such that w is a factor of $\mathbf{x}$ and z is a factor of $\mathbf{f}$. Let us assume that k is as small as possible. Since the Lyndon factors of length at most 3 in $\mathbf{x}$ and $\mathbf{f}$ coincide, we have $|w| > 3$, and thus $k \geq 5$. Recall that w can be written as $w = \lambda_w \mu_w$ where λ_w and μ_w are Lyndon words by Lemma 6. Furthermore, each of $|\lambda_w|$ and $|\mu_w|$ is a Fibonacci number, and therefore $|\lambda_w| + |\mu_w| = |w| = F_k$ yields $\{|\lambda_w|, |\mu_w|\} = \{F_{k-1}, F_{k-2}\}$. The same reasoning shows that $z = \lambda_z \mu_z$ and $\{|\lambda_z|, |\mu_z|\} = \{F_{k-1}, F_{k-2}\}$. But since $k \geq 5$, we have $F_{k-2} \geq 2$, which means that $\{\lambda_w, \mu_w\} = \{\lambda_z, \mu_z\}$ because the set of Lyndon factors of $\mathbf{x}$ of length less than F_k coincides with the set of Lyndon factors of $\mathbf{f}$ of length less than F_k and there is precisely one Lyndon factor of length F_{k-2} and precisely one Lyndon factor of length F_{k-1} . Consequently, z is a product of λ_w and μ_w . However, if $z = \lambda_w \mu_w$, then $z = w$, and if $z = \mu_w \lambda_w$, then z is not a Lyndon word, both contradictions. □

Remark 5. Theorem 4 shows that the shift orbit closure of a Fibonacci infinite word $\mathbf{f}$ is characterized by the mapping $\mathcal{L}_{\mathbf{f}}$, up to renaming letters. But in general, the mapping mapping $\mathcal{L}_{\mathbf{x}}$ does not determine a recurrent word $\mathbf{x}$. For example, the identity $\mathcal{L}_{\mathbf{x}} = \mathcal{L}_{\mathbf{y}}$

holds for the two periodic words $\mathbf{x} = (000001)^\omega$ and $\mathbf{y} = (000101)^\omega$, but clearly $F(\mathbf{x}) \neq F(\mathbf{y})$ and $F(\mathbf{x}) \neq F(c(\mathbf{y}))$, where c is the morphism $0 \mapsto 1, 1 \mapsto 0$.

5. ACKNOWLEDGEMENT

I thank Tero Harju for several useful discussions and for bringing my attention the result of Siromoney et al. (Lemma 12).

REFERENCES

- [1] J.-P. Allouche and J. Shallit. *Automatic Sequences: Theory, Applications, Generalizations*. Cambridge University Press, 2003.
- [2] J. Berstel and A. de Luca. Sturmian words, Lyndon words and trees. *Theoret. Comput. Sci.* **178** (1997), 171–203.
- [3] J. Cassaigne. On extremal properties of the Fibonacci word. *RAIRO-Theor. Inf. Appl.* **42**(4) (2008), 701–715.
- [4] M. G. Castelli, F. Mignosi, and A. Restivo. Fine and Wilf’s theorem for three periods and a generalization of Sturmian words. *Theoret. Comput. Sci.* **218** (1999), 83–94.
- [5] J.-M. Champarnaud, G. Hansel, and D. Perrin. Unavoidable sets of constant length. *International Journal of Algebra and Computation* **14**(2) (2004), 241–251.
- [6] K. T. Chen, R. H. Fox, and R. C. Lyndon. Free differential calculus, IV. The quotient groups of the lower central series. *Ann. Math.*, 2nd Ser., **68**(1) (1958), 81–95.
- [7] M. Crochemore and W. Rytter. Squares, cubes, and time-space efficient string searching. *Algorithmica* **13** (1995), 405–425.
- [8] J. D. Currie and K. Saari. Least periods of factors of infinite words. *RAIRO-Theor. Inf. Appl.* **43** (2009), 165–178.
- [9] A. de Luca. A combinatorial property of the Fibonacci words. *Inf. Process. Letters* **12**(4) (1981), 193–195.
- [10] V. Diekert, T. Harju, and D. Nowotka. Weinbaum factorizations of primitive words. *Russian Mathematics (Iz VUZ)* **54**(1) (2010), 16–25.
- [11] J.-P. Duval. Factorizing words over an ordered alphabet. *J. Algorithms* **4** (1983), 363–381.
- [12] A. Ehrenfeucht and D. M. Silberger. Periodicity and unbordered segments of words. *Discrete Math.* **26** (1979), 101–109.
- [13] S. Fischler. Palindromic prefixes and episturmian words. *J. Combin. Theory Ser. A* **113** (2006), 1281–1304.
- [14] N. J. Fine and H. S. Wilf. Uniqueness theorems for periodic functions. *Proc. Amer. Math. Soc.* **16**(1) (1965), 109–114.
- [15] H. Fredricksen and J. Maiorana. Necklaces of beads in k colors and k -ary de Bruijn sequences. *Discrete Math.* **23** (1978), 207–210.
- [16] D. E. Knuth. *The Art of Computer Programming*. Addison-Wesley, Reading, Mass., 1968.
- [17] M. Lothaire. *Combinatorics on Words*, Cambridge University Press, Cambridge, 1997.
- [18] M. Lothaire. *Algebraic Combinatorics on Words*, Cambridge University Press, Cambridge, 2002.
- [19] F. Mignosi, A. Restivo, and S. Salemi. Periodicity and the golden ratio. *Theoret. Comput. Sci.* **204** (1998), 153–167.
- [20] R. Siromoney, L. Mathew, V. R. Dare, and K. G. Subramanian. Infinite Lyndon Words. *Inf. Process. Letters* **50** (1994), 101–104.
- [21] R. Tijdeman and L. Zamboni. Fine and Wilf words for any periods II. *Theoret. Comput. Sci.* **410** (2009), 3027–3034.
- [22] Z.-X. Wen and Z.-Y. Wen. Some properties of the singular words of the Fibonacci word. *Europ. J. Combinatorics* **15** (1994), 587–598.

E-mail address: kasaar2@gmail.com